

Facial Rejuvenation Beauty Consult Intake Form

Doctor to review with Patient:

HISTORY

Skincare Products currently used: Cleanser, Exfoliant, Serum, SPF, Moisturizer Day/Night _____

Current Skincare includes: Retin-A, Hydroquinone, Kojic Acid, Glycolic Acid, Other _____

Which of the following aesthetic skincare services have you experienced in the past?

Facials, Microdermabrasion, Chemical Peels, Botox, Other _____

Laser, Intense Pulsed Light (IPL), Thermage, Cool Sculpt, Therma Cool, TCA, Other _____

As for the above two questions, were there any complications from the above previous procedures?

If yes, describe: _____

Do you feel faint with needles? Yes No

Have you ever been prescribed Acutane? Yes No

If yes, when? _____

Current medications and drugs, including topical _____

Are you pregnant or breastfeeding or do you plan to become pregnant in the near future? Yes No

Any known allergies: _____

Do you have history of allergic reaction to any of the anesthetic? Yes No

If YES, then specify which ones and when _____

Do you have any of the following: an autoimmune condition, a blood disorder, cancer, high blood pressure, acute allergies, herpes, cold sores, shingles, diabetes, contagious illness, headaches, high stress level, HIV, low or high blood pressure, a history of anesthetic reaction?

What areas of would you like to discuss? Hair, Face, Neck, Décolleté, Hands, Thighs, Buttocks, Stomach, Other _____

Please describe any concerns you have regarding your skin:


Please select the description that best describes your skin's sensitivity:

Little to no sensitivity

Mildly sensitive

Moderately sensitive

Very sensitive



Can cause *bruising*, could last 4-10 days, can be covered with makeup.

Can cause *facial redness*, which resembles a sunburn will last 6-12 hours or at the most 3 days; ice, cold compresses and a facial moisturizer can help to soothe the skin.

The numbing cream can cause a tingling sensation for some time (1-3 hours) after the injections.

Rarely causes infection, allergic reaction (i.e., to the numbing cream), bleeding during or after the procedure (do not take aspirin or anti-inflammatory medications for 7 days prior to the appointment), temporary injections site reactions such as *redness, pain/tenderness, firmness, swelling, lumps/bumps, bruising, itching & discoloration*.

NOTES: